



Genetic structure of the Painted Bunting and its implications for conservation of migratory populations

ANDREA CONTINA,^{1*}  JOSE L. ALCANTARA,² ELI S. BRIDGE,¹ JEREMY D. ROSS,¹ 
 WILLIAM F. OAKLEY,^{1,3} JEFFREY F. KELLY^{1,3,4} & KRISTEN C. RUEGG^{5,6} 

¹Oklahoma Biological Survey, University of Oklahoma, 111 East Chesapeake Street, SC Building 134, Norman, OK 73019, USA

²Colegio de Postgraduados, Km 36.5 Carr. Federal Mexico-Texcoco, Montecillo, Texcoco, Edo. De México 56230, Mexico

³Department of Biology, University of Oklahoma, 730 Van Vleet Oval, Norman, OK 73019, USA

⁴Corix Plains Institute, University of Oklahoma, Five Partners Place, 201 Stephenson Parkway, Norman, OK 73019, USA

⁵Department of Ecology and Evolutionary Biology, University of California Santa Cruz, Santa Cruz, CA 95064, USA

⁶Center for Tropical Research, Institute of the Environment and Sustainability, University of California, Los Angeles 619, Charles E. Young Dr. East, La Kretz Hall, Suite 300, Los Angeles, CA 90095, USA

The Painted Bunting *Passerina ciris* is a Neotropical songbird which breeds primarily in the USA during the summer and migrates to Mexico, Central America, southern Florida and the Caribbean over the winter. Male Painted Buntings are brightly coloured, which makes them highly sought after as pets, particularly in Mexico, Central America and Europe. We used short sequence repeats (microsatellite DNA) to investigate the population genetic structure of the Painted Bunting and its implications in conservation management of migratory populations. We found a detectable level of population differentiation as revealed by pairwise F_{ST} and R_{ST} comparisons and Bayesian clustering analyses, with strong support for differentiation between eastern and western Painted Buntings (e.g. Oklahoma and Georgia $F_{ST} = 0.1$; $P = 0.005$; $R_{ST} = 0.18$; $P = 0.04$) in accordance with previous mitochondrial DNA analysis. We recovered additional support for two subgroups within the western clade. While linking migrant songbirds captured outside of the USA to their breeding populations remains a challenge, we show that natural levels of population genetic differentiation can be detected via microsatellite DNA markers and exploited in migratory connectivity studies. We also demonstrate the potential utility of our low-cost markers for population identification of birds recovered from the pet trade by screening a small subset of samples ($n = 5$) collected as part of wildlife tracking. We discuss the implications of our results for future efforts to understand patterns of population decline in Painted Buntings more generally, as well as how we might expand this methodology to combat illegal pet-trade activity in this and other songbird species.

Keywords: Convention on International Trade in Endangered Species of Wild Fauna and Flora, forensics, international pet trade, migration, population genetics.

Over half of the Neotropical migrant bird species found breeding in North America have shown marked declines in abundance over the last several

decades (Robbins *et al.* 1989, Sauer *et al.* 2013). Population declines are thought to relate to stressors encountered by migrants at each stage in the annual cycle, including habitat loss, predation by house cats, exposure to disease, illegal poaching and global climate change (Jonzén *et al.* 2006,

*Corresponding author.
 Email: andrea.contina@ou.edu
 Twitter: @andrea_contina




Figure 1. The Painted Bunting annual species distribution: the summer distribution is highlighted in yellow (breeding) and wintering distribution (c. September to April) is marked in dark blue. Red triangles indicate sampling locations within the natural species distribution range. Black triangles indicate market sampling locations in Mexico and Italy. Map source BirdLife International. [Colour figure can be viewed at wileyonlinelibrary.com]

Altizer *et al.* 2011, Loss *et al.* 2013). For migrants, ecologists have long recognized that conservation measures on the breeding grounds are most effective when wintering and migratory ecology are understood, and winter and migratory stopover habitat protected (Sherry & Holmes 1993, Marra *et al.* 1998, Norris & Taylor 2006), but such information is often difficult to attain.

The Painted Bunting *Passerina ciris* is a neotropical migratory songbird that is estimated to have declined by ~ 55% in some parts of its range over the last 30 years (Sauer *et al.* 2003, 2013). This species breeds in a variety of habitats across the southern USA and northern Mexico, wintering in parts of Mexico, Central America, the Florida Keys and Florida Peninsula, and the Bahamas (Rappole & Warner 1980, Howell & Webb 1995, Lowther

et al. 1999, Mlodinow & Hamilton 2005, Bridge *et al.* 2011). While early taxonomic studies identified two subspecies within Painted Buntings (*P. c. pallidior* and *P. c. ciris*) separated by a zone of introgression in the central USA (Storer 1951), phylogenetic analysis using mitochondrial DNA supports the idea that the major split between the two subspecies is defined by a 550-km gap between eastern Mississippi and western Georgia (Herr *et al.* 2011, see also Thompson 1991a,b, Shipley *et al.* 2013) (Fig. 1). Breeding Bird Survey data suggest that the eastern population in particular is declining and that several populations from the south-central and eastern United States and northeastern Mexico are now locally extirpated from parts of their former range (Sauer *et al.* 2003, NAS 2004a,b, USFWS 2004). As a result,

the Painted Bunting has been listed as Near Threatened on the IUCN Red List, as a species of conservation concern by both the U.S. Fish and Wildlife Service and Partners in Flight, and as Near Threatened by Birdlife International (Sauer *et al.* 1997, Sykes & Holzman 2005, Sykes *et al.* 2007, USFWS 2008).

Habitat loss (both at wintering and at stopover sites), habitat fragmentation and brood parasitism by the Brown-headed Cowbird *Molothrus ater* may all contribute to declines in Painted Buntings, but it has also been suggested that capture for the international cage bird trade is also a particularly serious threat (Hamilton 2001, Iñigo-Elias *et al.* 2002, Lopez Medellin & Iñigo-Elias 2002, Arizmendi Arriaga & Ramos Rivera 2017). According to some historical estimates, nearly 15 000 Painted Buntings were collected and exported every year from Mexico from the mid-1970s up until the early 1980s (Iñigo-Elias 1986). In 1982, Mexico banned wildlife exports, but over a decade later, Mexico resumed the international trade of wild animals, mostly to South America, northern Europe and Malaysia (Ramos 1982, Iñigo-Elias *et al.* 2002). Since 1999, Mexico has utilized capture quotas and management unit areas (UMAs) to promote and monitor sustainable harvesting of Painted Buntings and other wildlife, but precise control of these quotas is difficult to enforce (SEMARNAT 2013). In 2004, the Painted Bunting was proposed for inclusion in the Convention on International Trade in Endangered Species of Wild Fauna and Flora (CITES, Appendix II), but it did not meet listing criteria (under Resolution Conf. 9.24) because the official annual harvested and exported stock from Mexico was < 0.3% of the estimated global population size (IUCN and TRAFFIC 2004). The lack of CITES protection makes it extremely difficult to quantify legal and illegal captures, as well as international exports of this threatened species (Cruz-Romo & de Oliveras Ita 2011).

One option for understanding the role that pet trafficking and other anthropogenic stressors might play in population declines in the Painted Bunting is to use genetic markers to identify samples collected anywhere along the migratory pathway back to their breeding population of origin. Modern molecular markers such as mitochondrial DNA, short sequence repeats (microsatellites) and single nucleotide polymorphisms (SNPs) are useful intrinsic markers for determining population

structures as well as for tracking large-scale movements of migratory individuals (Aulsebrook 2004, 2008, Rundel *et al.* 2013, Ruegg *et al.* 2014, Contina *et al.* 2017). The same sorts of genetic tools have been employed in a forensic DNA framework to identify individuals that have been confiscated as a result of illegal wildlife trafficking (Iyengar 2014). While these forensic approaches have been used effectively on several species such as baleen whales, fish, macaws and lizards (Baker & Palumbi 1994, Ogden *et al.* 2009, Galimberti *et al.* 2013, Welton *et al.* 2013a, Presti *et al.* 2015), to date the efficacy of such techniques has not been tested in an exploited neotropical migratory bird system.

Here we investigate patterns of population structure in Painted Buntings from across the species breeding and wintering range using microsatellite DNA markers. We begin with an assessment of population structure across the breeding range and discuss the utility of our markers for migratory connectivity analysis more generally. We then take the opportunity to test the utility of our markers on a small subset of samples from the pet trade and discuss the implications of our results in the Painted Bunting as well as the use of genetic markers for combating wildlife trafficking more broadly.

METHODS

Natural population dataset and pet-trade examples

Our genetic sampling comprised 131 Painted Buntings sampled at 15 sites across the species breeding and wintering geographical range (Fig. 1). Our dataset included DNA aliquots for 50 birds previously sequenced by Herr *et al.* (2011) and purified genomic DNA for 81 adult birds using blood, muscle and feather samples from museum collections and mist-netting operations (Supporting Information Table S1).

We included samples from two distant populations in Louisiana – Johnson Bayou and Bayou Cocodrie – which are about 245 km apart. However, our field site at Bayou Cocodrie was practically contiguous to the Mississippi state border; therefore, we refer to the samples collected in Bayou Cocodrie as from a ‘Mississippi population’ for a better geographical characterization and to distinguish them from the samples collected at Johnson Bayou in southwestern Louisiana. At the wintering grounds in Costa Rica, eight birds were

captured at sites < 40 km apart (Tamarindo and Guardia), so we treated them as if they were obtained from a single population.

To provide an example of an applied pet-trade forensic investigation, we acquired five Painted Bunting samples from the pet market in central Mexico and Europe, and performed genetic assignment tests. In Mexico, between October and December 2013, we visited eight markets in Mexico City that were referred to us for trading in birds. We found Painted Buntings for sale in three markets for about 10 USD each, and tail feathers were taken from four individuals: one adult male was acquired from Mercado La Argentina; one bird of unknown sex and age was acquired from Mercado de Xochimilco; and one adult male together with one bird of unknown sex and age were acquired at the Catedral de Texcoco, a town about 20 km northeast of Mexico City. We relied on plumage characteristics to age and sex each bird according to Pyle (1997). In Italy, we acquired one male Painted Bunting from a large distributor of exotic birds located near Pavia for 700 USD.

DNA extraction

We stored ~30 μ L of whole blood from the brachial vein in 0.4 mL of Queen's lysis buffer at 4 °C (Seutin *et al.* 1991). Muscle tissues from Sinaloa birds was frozen in liquid nitrogen and stored at -20 °C. We stored primary and tail feathers (first or ninth primary feather or first and second rectrices) at room temperature in paper envelopes to absorb humidity and allow the feather calamus to dry, thus limiting DNA degradation. We purified DNA from blood and muscle samples using the Qiagen DNA extraction kit (Qiagen Inc., Valencia, CA, USA) according to the protocols supplied by the manufacturer, whereas DNA from feather samples was purified following a modified feather digestion procedure (8 h at 57 °C) using a DNA extraction mixture without dithiothreitol (modified from De Volo *et al.* 2008).

Microsatellite genotyping

We made use of 13 microsatellite primer pairs developed by Contina *et al.* (2016a,b) (Supporting Information Table S2) to PCR-amplify loci, and visualized the electropherograms and scored allele

sizes in PEAKSCANNER 2.0 (Applied Biosystems, Carlsbad, CA, USA). We implemented a microsatellite genotyping method following the M13 hybrid primer technique to reduce the costs of molecular laboratory work (Schuelke 2000). We analysed each DNA sequence using an ABI 3130XL sequencer with an internal size standard (Genescan LIZ-600; Applied Biosystems). We performed PCR following thermocycler conditions detailed in Contina *et al.* (2016a). We used MICRO-CHECKER (Van Oosterhout *et al.* 2004) to detect the presence of null alleles, large allele dropout and scoring errors. We did not rescore individuals to estimate an error rate.

Population genetic diversity and differentiation

We used ARLEQUIN 3.5.1.3 (Excoffier & Lischer 2010) to compute several population genetic tests implementing the Markov chain Monte Carlo (MCMC) algorithm using 10^6 iterations, 10^5 dememorization steps and significance level $\alpha < 0.05$. First, we computed the expected and observed heterozygosity for each locus within each population and tested for deviation from Hardy-Weinberg equilibrium (HWE) (Guo & Thompson 1992) and linkage disequilibrium (LD) (Slatkin & Excoffier 1996). Then, we compared populations by computing pairwise F_{ST} and R_{ST} distances (Weir & Cockerham 1984, Raymond & Rousset 1995, Slatkin 1995). We used the program FSTAT (Goudet 2001), which corrects for variation in the number of sampled individuals, to calculate the number of alleles and allelic richness per locus and population to make comparisons across the species range (Nei 1988, Petit *et al.* 1998). We tested the isolation-by-distance (IBD) model (Wright 1943) in GENEPOP through the ISOLDE option, which runs a regression of $F_{ST}/(1 - F_{ST})$ estimates for population pairs on the logarithm of their geographical distances (Raymond & Rousset 1997).

Population genetic structure

We ran the Bayesian clustering algorithm implemented in the program STRUCTURE 2.3.2.1 (Pritchard *et al.* 2000) to investigate the potential for population structure across the breeding grounds with the full set of 13 Painted Bunting loci developed by Contina *et al.* (2016a). We ran STRUCTURE using the correlated allele

frequencies prior and both with and without the location prior function (LOCPRIOR). When using the LOCPRIOR, individuals sampled from the same area are assumed to have their admixture proportions (Q values) drawn from the same Dirichlet distribution (Hubisz *et al.* 2009). We tested the number of clusters $K = 1-6$ using 10 replicate simulations of 10^5 MCMC repetitions each and a burn-in of 10^5 iterations. We determined the most likely value of K by following the ΔK method (Evanno *et al.* 2005) implemented in STRUCTURE HARVESTER (Earl & vonHoldt 2012). We merged the results for the best model with the highest likelihood assignment scores (Francois & Durand 2010) with CLUMPP (Jakobsson & Rosenberg 2007), and converted the output into a postscript file with DISTRUCT (Rosenberg 2004). We visualized the postscript file in Adobe ILLUSTRATOR CC (Adobe Systems, San Jose, CA, USA).

Assignment of wintering and pet-trade individuals

To assign birds of unknown origin back to their breeding population we used the USEPOPINFO function within a population admixture framework in STRUCTURE. We used the result of our population structure analysis from above to inform the separation of the breeding grounds into three separate groups: (1) western breeders (Central TX and OK), (2) central breeders (Western TX, Eastern TX, AR, LA, MS) and (3) eastern breeders (GA and NC). Samples from the wintering grounds (Sin, Oax, Yuc, GU, NI, CR) and our five pet-trade individuals were coded as 'migrants from unknown breeding populations' and assigned to one of the three breeding populations using 10 replicate simulations of 10^5 MCMC repetitions each and a burn-in of 10^5 iterations, and we used the default parameters GENSBACK = 2 and MIGRPRIOR = 0.05.

We also ran a genotype assignment test based on allele frequencies for samples collected in Oklahoma, a representative western breeding population, as well as samples collected in North Carolina, a representative eastern breeding population, and used the pairwise log-likelihood assignment probabilities calculated in ARLEQUIN to visualize the extent of differentiation between these populations following Paetkau *et al.* (1997) and Waser and Strobeck (1998).

RESULTS

Population genetic diversity and structure

We provide insight on the genetic diversity of Painted Bunting populations across most of the species breeding and wintering range in the USA and Central America through PCR-amplification of 3406 alleles across 13 microsatellite loci among 131 individuals (Dryad Digital Repository: <https://doi.org/10.5061/dryad.mg4nc21>). A detailed account of the number of alleles sampled and allelic richness per locus and population across the species distribution range is presented in Table 1. No sample failed to PCR-amplify at any locus and we did not find evidence for large allele dropout or scoring error due to stuttering.

We found significant pairwise F_{ST} and R_{ST} distances in 71.4% and 24.7% of all comparisons across 15 populations, respectively, with larger values generally observed between populations of more distant sites (Table 2, Supporting Information Table S6). The test for isolation-by-distance (IBD) recovered a moderate but significant correlation between genetic differentiation (F_{ST}) and geographical distance ($R^2 = 0.46$; $P = 0.005$). F_{ST} values ranged from 0.008 to 0.12 whereas R_{ST} values ranged from 0.001 to 0.31 across all the populations from the breeding and wintering grounds. The highest significant F_{ST} value was between Georgia and Costa Rica ($F_{ST} = 0.12$, $P = 0.02$) and the lowest was between Arkansas and eastern Texas ($F_{ST} = 0.008$, $P = 0.48$). The highest significant R_{ST} value was between Oklahoma and western Texas ($R_{ST} = 0.31$, $P < 0.0001$) and the lowest, but significant, value was between Oklahoma and Sinaloa ($R_{ST} = 0.04$, $P = 0.02$). We also recovered a very low F_{ST} value between Oklahoma and Sinaloa ($F_{ST} = 0.01$, $P = 0.07$), the latter of which is known to host birds migrating from Oklahoma in the autumn (Contina *et al.* 2013). We found that locus 11 (Pb11 in Table S2) had the highest allelic richness value (ARV) across populations (ARV locus 11 = 3.6). Some populations showed loci deviating from Hardy-Weinberg equilibrium (HWE) and evidence of linkage disequilibrium (Supporting Information Tables S3 & S4). As results for HWE and LD were not consistent among populations, we use all the loci in the analyses reported below.

STRUCTURE suggested differentiation among samples from the western, central and eastern

Table 1. (a) Number of alleles sampled per locus and population. (b) Allelic richness per locus and population. Population sample size is provided in parentheses.

	Central Texas (2)	Oklahoma (21)	Western Texas (6)	Eastern Texas (6)	Arkansas (8)	Louisiana (8)	Mississippi (8)	Georgia (2)	North Carolina (12)	Sinaloa (15)	Oaxaca (7)	Guatemala (6)	Nicaragua (8)	Costa Rica (8)	Yucatan (14)	
(a)																
Number of alleles																
Locus 1	3	6	4	4	5	5	4	1	2	10	4	3	5	5	6	6
Locus 2	3	10	4	4	4	5	3	1	4	7	7	4	6	6	6	6
Locus 3	4	18	10	6	8	8	10	3	11	16	7	8	9	9	12	12
Locus 4	4	11	6	4	5	8	6	1	4	7	9	6	9	9	7	7
Locus 5	3	14	9	3	7	8	6	2	6	14	7	6	5	10	12	12
Locus 6	2	8	3	4	5	8	10	4	7	10	3	8	4	4	8	8
Locus 7	2	13	6	8	9	10	8	4	8	13	7	7	6	5	10	10
Locus 8	2	4	3	4	5	3	4	3	5	5	3	3	4	3	2	2
Locus 9	3	12	7	8	8	8	7	4	11	15	9	7	9	5	9	9
Locus 10	3	8	3	4	3	2	6	1	8	7	5	3	7	4	4	4
Locus 11	3	25	9	9	11	14	13	3	9	17	10	9	12	9	18	18
Locus 12	4	7	5	5	5	3	5	4	6	10	4	3	5	5	6	6
Locus 13	2	9	6	7	10	9	8	4	9	10	5	7	8	8	11	11
(b)																
Allelic richness																
Locus 1	3	2.599	2.925	2	2.719	1.999	2.409	1	1.544	2.933	2.432	2.149	2.676	2.975	2.606	2.606
Locus 2	3	2.689	2.624	2.624	2.48	2.81	2.497	1	2.276	3.016	3.384	2.697	2.859	3.104	2.789	2.789
Locus 3	4	3.723	3.818	3.008	3.435	3.514	3.614	3	3.427	3.638	3.374	3.564	3.485	3.607	3.55	3.55
Locus 4	4	2.985	3.051	2.495	2.699	3.283	2.97	1	2.44	2.67	3.56	3.18	3.485	3.419	2.846	2.846
Locus 5	3	3.292	3.727	1.667	2.988	3.038	3.092	2	2.602	3.272	3.193	3.051	2.197	3.181	3.341	3.341
Locus 6	2	2.931	2.404	2.697	2.719	3.038	3.657	4	3.106	3.169	2.181	3.564	2.626	2.681	2.601	2.601
Locus 7	2	3.104	3.382	3.564	3.564	3.614	3.326	4	3.136	2.986	3.193	3.172	2.769	2.865	3.254	3.254
Locus 8	2	1.974	1.909	2.495	2.81	2.142	2.791	3	2.924	1.971	1.791	2.434	2.445	1.975	1.76	1.76
Locus 9	3	3.408	3.473	3.491	3.392	3.326	3.142	4	3.49	3.588	3.515	3.343	3.485	2.809	3.155	3.155
Locus 10	3	2.609	2.182	2.402	2.312	1.816	2.97	1	3.356	2.474	2.712	2.404	3.031	2.147	2.383	2.383
Locus 11	3	3.823	3.655	3.727	3.707	3.9	3.807	3	3.392	3.597	3.681	3.727	3.678	3.376	3.874	3.874
Locus 12	4	2.645	2.917	3.145	2.699	2.115	2.92	4	2.775	3.364	2.591	2.311	2.81	2.845	2.838	2.838
Locus 13	2	2.812	3.008	3.271	3.578	3.285	3.356	4	3.378	3.215	2.961	3.271	3.356	3.313	3.56	3.56

Table 2. F_{ST} values for pairwise comparisons of populations calculated in ARLEQUIN (lower diagonal). Significant values ($P < 0.05$) are highlighted in bold. Pairwise population geographical distances (km) are indicated in italics along the top diagonal.

F_{ST}	Central		Western		Eastern		North				Costa			
	Texas	Oklahoma	Texas	Oklahoma	Texas	Oklahoma	Georgia	Carolina	Sinaloa	Oaxaca	Guatemala	Nicaragua	Rica	Yucatan
Central		312	458	404	562	956	739	1670	1967	1097	1635	2060	2638	1578
Texas														
Oklahoma	0.011		696	471	470	993	758	1626	1866	1343	1939	2335	2872	1778
Western	0.038	0.05		830	1020	1340	1150	2093	2406	642	1545	2063	2719	1790
Texas														
Eastern	0.072	0.073	0.047		257	561	335	1269	1573	1417	1605	1917	2431	1302
Texas														
Arkansas	0.045	0.049	0.014	0.008		551	311	1153	1418	1628	1842	2117	2577	1440
Louisiana	0.106	0.088	0.047	0.01	0.018		245	765	1113	1881	1637	1745	2111	2248
Mississippi	0.077	0.067	0.05	0.035	0.009	0.024		941	1265	1723	1665	1887	2293	2424
Georgia	0.142	0.104	0.116	0.065	0.078	0.088	0.073		403	2629	2218	2157	2276	1275
North	0.096	0.086	0.086	0.05	0.056	0.06	0.031	0.017		2985	2611	2547	2606	1667
Carolina														
Sinaloa	0.023	0.012	0.052	0.08	0.051	0.094	0.071	0.099	0.1		1451	2540	2745	2046
Oaxaca	0.037	0.018	0.04	0.048	0.041	0.082	0.06	0.111	0.083	0.035		615	1325	1395
Guatemala	0.018	0.035	0.042	0.021	0.01	0.045	0.021	0.052	0.029	0.051	0.039		690	775
Nicaragua	0.043	0.035	0.039	0.046	0.035	0.072	0.032	0.093	0.039	0.06	0.028	0.034		144
Costa Rica	0.076	0.073	0.081	0.048	0.044	0.065	0.06	0.126	0.089	0.096	0.059	0.065	0.03	1275
Yucatan	0.096	0.092	0.084	0.081	0.05	0.059	0.062	0.1	0.088	0.101	0.097	0.06	0.1	0.058




Figure 2. Individual Q matrix calculated in STRUCTURE. The graph shows the assignment probabilities (Q) of Painted Bunting clusters across the species breeding distribution range. Each area within two vertical black lines represents a population and the area width is proportional to the number of individuals sampled. The probability (Q) of each individual to be assigned to one or more clusters is shown on the vertical axis and the clusters are represented by different colours. The figure shows results for $K = 3$ based on the admixture (ADMX) and admixture + LOCPRIOR (ADMX + LOCPRIOR) models, respectively. The two models yielded similar results with slightly less evident population structure when LOCPRIOR was not implemented. [Colour figure can be viewed at wileyonlinelibrary.com]

breeding populations (Fig. 2, Supporting Information Fig. S2). We found that the most likely hierarchical group (K) was equal to two for the admixture model according to the Evanno *et al.* (2005) ΔK method and three when we included the location information as a prior (LOCPRIOR; Supporting Information Fig. S1). The Bayesian population structure analysis based on either the admixture or the admixture + LOCPRIOR model suggested a similar sub-division within the species range for higher values of K (e.g. $K \geq 3$; Fig. S2).

Overall, the population subdivision at the breeding grounds supported evidence for three main groups: a western cluster covering Oklahoma and central Texas, a central cluster formed by eastern and western Texas, Arkansas, Louisiana and Mississippi, and an eastern group that includes the two breeding populations along the Atlantic coast of the USA (Fig. 2 and Fig. S2). STRUCTURE identified some individuals, possibly migrants from the central cluster, within the western and eastern breeding populations as indicated by the red Q -bars in Figure 2 (see also Fig. 3 USEPOPINFO results for further details). The breeding populations of the central cluster shared some genetic

similarity with individuals sampled from the eastern cluster (Georgia and North Carolina; Fig. 2).

Origin of migrants wintering in Central America

Although we could not determine the breeding origin of all migrants, many birds sampled on the wintering grounds were confidently assigned to one of the three putative breeding region clusters (western, central, eastern; Fig. 3). The lack of assignment power might be due to admixed birds breeding between our two sampled breeding source populations or, alternatively, to the occurrence of unsampled genetic variation on the breeding grounds, which, if included, would improve assignments of wintering birds. About 60% of migrants that passed through Sinaloa and to some extent Oaxaca could be assigned to western breeders with confidence ($Q > 0.8$). Migrants wintering in Guatemala, Nicaragua, Costa Rica and the Yucatan Peninsula showed a mixture of assignment probabilities to one of the three breeding groups (e.g. individuals 98, 107, 116 and 125 in Fig. 3 and Supporting Information Table S5), perhaps reflecting gaps in our sampling




Figure 3. Individual Q matrix calculated in STRUCTURE within a population admixture framework and the function USEPOPINFO for three breeding clusters. Painted Bunting samples from populations occurring in the western, central and eastern part of the breeding range are pooled together and colour-coded as blue, red and yellow, respectively, in accordance with STRUCTURE model results (see Fig. 2 and Supporting Information) to test for the origin of wintering migrants and birds sold in the international pet market. Migrants sampled at a moulting site in northwestern Mexico (Sinaloa), and to some extent at a wintering site in southern Mexico (Oaxaca), show strong genetic similarities with western breeders. The proportion of ‘western’, ‘central’ and ‘eastern’ genetic signatures in pet trade birds is strongly skewed toward the ‘central’ (red) cluster. [Colour figure can be viewed at wileyonlinelibrary.com]

effort across the breeding grounds. A few migrants wintering in the Yucatan Peninsula could be assigned to eastern breeders ($Q > 0.6$; e.g. individuals 118 and 122 in Fig. 3 and Supporting Information Table S5), whereas pet-trade samples were consistently associated (Q scores ranged from 0.6 to 0.77) with the breeding birds of the central clade (Fig. 3, Table S5).

To illustrate the resolving power of the genotype analysis based on population allele frequencies computed in ARLEQUIN, we plotted the pairwise log-likelihood assignment probabilities of individuals from Oklahoma (western breeding population) against the log-likelihood assignment probabilities of individuals from North Carolina (eastern breeding populations). This plot shows marked differences in assignment probabilities between the representative samples from the western and eastern population, although no apparent plumage or other morphological differences are evident (Fig. 4).

DISCUSSION

Our research demonstrates that Painted Buntings can be separated into three main groups on the

breeding grounds using microsatellite markers – a western, central and eastern group. By taking advantage of the genetic distinctiveness on the breeding grounds, we demonstrate how microsatellites can serve as a low-cost method (about 10 USD per sample) for identifying the population of origin of birds captured from anywhere along their migratory cycle, as well as sold in the international avian pet market. Below we outline the implications of our findings for our understanding of migratory connectivity in the Painted Bunting, as well as the potential use of microsatellite markers for forensic analysis in this species.

Overall, we recovered F_{ST} and R_{ST} distances between pairwise comparisons of populations that followed a pattern of differentiation in agreement with the STRUCTURE results, with greater differentiation between more distant sampling sites. However, in the STRUCTURE results we also recovered a signature of eastern alleles occurring within the central clade, whereas the western clade appeared separated from the central group despite geographical continuity (Figs 1 & 2). Storer (1951) studied Painted Bunting morphology and plumage coloration and claimed that the




Figure 4. Log-likelihood probability values of western birds (Oklahoma) being assigned back to their breeding population (blue circles) plotted against the log-likelihood probability values of being assigned to the North Carolina breeding population (yellow squares). (a, b) Representative adult male Painted Bunting photographed in Oklahoma (western population) and South Carolina (eastern population), respectively. The two individuals are practically indistinguishable to the human eye. However, the genetic assignment test can help to differentiate individuals from different populations. Photo credits: Crystina Meyers (a) and William Oakley (b). [Colour figure can be viewed at wileyonlinelibrary.com]

two sub-species met along a zone of introgression running through the middle of the western range, providing an expanded range to '*P. c. ciris*' to the west and ignoring the species' discontinuous geographical breeding distribution. The concept of two breeding populations of Painted Bunting connected through a zone of introgression running across parts of Oklahoma and Texas was supported by the results from STRUCTURE analyses without the LOCPRIOR function implemented (e.g. $K = 2$ in Fig. S2). Furthermore, Battey *et al.* (2017) investigated the possibility of a migratory divide in the Painted Bunting using single nucleotide polymorphisms (SNPs) and showed evidence of a zone of introgression running through the western part of the species breeding range. Our microsatellite DNA results also support the occurrence of genetic differentiation among populations within the western breeding range (Fig. 2 and Fig. S2). The biogeographical underpinning of this pattern remains difficult to explain even in light of the results provided by Battey *et al.* (2017). We speculate that differences in moult and migration strategy have evolved across the species range in response to a variety of environmental conditions. Determining where the zone of introgression occurs between the western and central breeding clades will improve our understanding of

which factors contributed to the population differentiation patterns described here and elsewhere.

Furthermore, when we implemented the LOCPRIOR function in STRUCTURE we found clear support for three populations on the breeding grounds (Fig. 2 and Fig. S1). Because one of the objectives of the present study was to test the utility of microsatellite markers for population assignment of non-breeding birds to breeding locations at as fine a spatial scale resolution as possible, we used $K = 3$ as the number of source populations on the breeding grounds and added usable information (i.e. LOCPRIOR) to our assignment tests.

Our results suggest that microsatellite DNA can be successfully adopted in the Painted Bunting to study population genetic structure, assign breeding origin of migrants or pet-trade birds, and in some cases investigate sequence repeats in candidate genes related to migration (Contina *et al.* 2016b, 2018). On the other hand, the full potential of adopting an SNP approach might emerge once genomic variant screening and selection are conducted through samples collected from multiple breeding populations in proximity of the zone of introgression, where an informative sub-set of variants is developed through reproducible molecular assays, and novel candidate genes under selection

are discovered throughout the genome (DePristo *et al.* 2011, Jonker *et al.* 2012, Bay *et al.* 2018).

Through our analysis of assignment posterior probabilities using the program STRUCTURE, we found that birds wintering in Guatemala, Nicaragua, Costa Rica and the Yucatan Peninsula were most likely from the central and eastern breeding population, whereas birds sampled from Sinaloa and, to some extent, Oaxaca (southern Mexico) were most likely from the western clade. However, our assignment probabilities for 22 wintering samples (about 35% of our migratory bird dataset) did not resolve their breeding origin with confidence (grey colours in Table S5). This may result from unsampled genetic variation on the breeding grounds or because the central population represents a transition zone between the western and eastern group (see also Battey *et al.* 2017).

Our assignment results for the Sinaloa and about half of the Oaxaca samples are consistent with the pattern for the western population recently inferred by Contina *et al.* (2013). Contina *et al.* (2013) implemented an ultra-light geolocator design to track Painted Buntings from a breeding population in Oklahoma and revealed a moult-migration strategy: birds moving westward at the end of the summer toward Sinaloa (northwest Mexico) stopped to moult before continuing along the west coast of Mexico and then returned to the breeding sites in the spring along the east coast of Mexico. It is thought that this moult-migration strategy may allow Painted Buntings to exploit higher biological productivity resulting from monsoonal rains in Mexico (Rohwer *et al.* 2005, Pyle *et al.* 2009, Bridge *et al.* 2016). Laws for the conservation of migratory species take into account spatial and temporal challenges of protecting populations that move across natural landscapes and geopolitical borders (Ellison 2014) and we recommend that conservation action plans should specifically address the migratory behaviour occurring across the entire annual cycle (Marra *et al.* 2015). Therefore, we propose that Sinaloa should be considered a target area for prioritizing conservation efforts for the Painted Bunting and potentially other migrant species.

Although wintering population contact was previously considered unlikely on the basis of an earlier study which examined differences in wing-length across wintering locations and used such morphological differences to assign birds to breeding populations of origin (Thompson 1991a), our

results suggest possible contacts between individuals from the western and eastern populations on the wintering grounds (e.g. Yucatan Peninsula). Our results are thus consistent with banding records that indicate that at least some eastern Painted Buntings migrate through Cuba and the Caribbean (Sykes *et al.* 2007). Our findings are important in light of conservation management of separate breeding populations of Painted Bunting, including the small and declining eastern bunting population, as wintering locations hosting a mixture of western and eastern buntings should be closely monitored as potential sites for conservation efforts.

In addition to using our markers to identify patterns of migratory connectivity across the range, we were also able to identify the population of origin of a small sub-sample of birds sold in the international avian pet market (both Mexico and Europe). This exploratory examination showed that the pet-market birds are most likely from the central breeding population of Painted Buntings (Fig. 3, Table S5). Due to our small sample size we do not exclude the possibility that many buntings sold as pets worldwide may have been sourced from populations breeding along the Atlantic coast of the USA and wintering in Central America and the Caribbean. We simply note that our low-cost molecular approach to assign pet-market samples to natural populations is a valid method with great investigative potential in forensic genetic analyses of the Painted Bunting and other songbirds exploited in the international pet-trade industry, such as Indigo Bunting *Passerina cyanea*, Lazuli Bunting *Passerina amoena*, Blue Grosbeak *Passerina caerulea* and Northern Cardinal *Cardinalis cardinalis*. We note that microsatellite analyses presently offer a cheaper and easier option compared with many genomic approaches, as they can be easily implemented on degraded and low concentrations of DNA and are easier to add new samples to than, for example, ddRAD approaches such as those used by Battey *et al.* (2017) for the Painted Bunting.

CONCLUSIONS

Our findings highlight the importance of songbird migration stopover sites and wintering areas in relation to conservation plans for multiple breeding populations coming into contact during specific phases of their annual cycle. The genetic

distinctiveness and negative population trend of the eastern Painted Bunting populations (e.g. Georgia, North Carolina) supports the idea that this population represents an evolutionarily significant unit (ESU, *sensu* Moritz 1994) in agreement with Herr *et al.* (2011). The demonstrated utility of our markers for connectivity and forensic analysis suggests that microsatellites may function as a low-cost alternative for population identification in the Painted Bunting and other closely related migratory species.

We thank Connie Herr, Paul Sykes and John Klicka for providing 50 Painted Bunting DNA aliquots used in this study and Sievert Rohwer for collecting tissue samples in Sinaloa. We thank Ann Harris and the Core Molecular Biology Laboratory at the University of Oklahoma for helping with laboratory work. Larry Weider, Michael Paten, Bruce Hoagland, Tom Smith, two reviewers, Jon Martin Collinson and Rauri Bowie provided valuable comments on earlier drafts of the manuscript. We are grateful to Tyler Michels, Alice Andolfatto and the many field assistants who helped collect samples across the USA and Europe. We thank The Institute for Bird Populations together with MAPS and MoSI banding station managers, and UCLA – Center for Tropical Research, for providing feather samples. We thank the Wichita Mountains Wildlife Refuge in Oklahoma and St. Catherine Creek National Wildlife Refuge in Mississippi for allowing us to work in special use areas. We thank Frank Moore, Kristen Covino and the Migratory Bird Research Group of the Department of Biological Sciences at The University of Southern Mississippi in Hattiesburg for collecting bunting samples at Johnson Bayou, Louisiana. All work with animals was performed with relevant state and federal permits (#23215) and was approved by the Institutional Animal Care and Use Committee of the University of Oklahoma. This research was funded by the National Science Foundation (IDBR 1014891, DGE 1545261 and DEB 0946685) and by the United States Department of Agriculture (NIFA-AFRI-003536). Additional funding for covering the costs of ‘open access’ publication was provided by the University Strategic Organization in ‘Applied Aeroecology’ at the University of Oklahoma.

REFERENCES

- Altizer, S., Bartel, R. & Han, B.A. 2011. Animal migration and infectious disease risk. *Science* **331**: 296–302.
- Arizmendi Arriaga, M.D.C. & Ramos Rivera, P. 2017. *Evaluación del estado de conservación de las poblaciones de Passerina ciris, P. leclancherii, P. amoena, P. rositae, P. caerulea, Cardinalis cardinalis y C. sinuatus en México: recomendaciones para su conservación y aprovechamiento sustentable*. Version 1.3. Comisión nacional para el conocimiento y uso de la biodiversidad. Occurrence dataset <https://doi.org/10.15468/qdmngel> accessed via GBIF.org on 2018-07-28.
- Avise, J.C. 2004. Chapter 9: Molecular Markers in Conservation Genetics. In *Molecular markers, natural history, and evolution* (2nd ed.): 495–500. Sunderland, MA: Sinauer Associates.
- Avise, J.C. 2008. The history, purview, and future of conservation genetics. In Carroll, S.P. & Fox, C.W. (eds) *Conservation Biology: Evolution in Action*: 5–15. Oxford: Oxford University Press.
- Baker, C.S. & Palumbi, R.S. 1994. Which whales are hunted? Molecular genetic evidence for illegal whaling. *Science* **265**: 1538–1539.
- Batley, C.J., Linck, E.B., Epperly, K.L., French, C., Slager, D.L., Sykes, P.W. & Klicka, J. 2017. A migratory divide in the Painted Bunting (*Passerina ciris*). *Am. Nat.* **191**: 259–268.
- Bay, R.A., Harrigan, R.J., Underwood, V.L., Gibbs, H.L., Smith, T.B. & Ruegg, K. 2018. Genomic signals of selection predict climate-driven population declines in a migratory bird. *Science* **359**: 83–86.
- Bridge, E.S., Fudickar, A.M., Kelly, J.F., Contina, A. & Rohwer, S. 2011. Causes of bimodal stable isotope signatures in the feathers of a molt-migrant songbird. *Can. J. Zool.* **89**: 951–959.
- Bridge, E.S., Ross, J.D., Contina, A.J. & Kelly, J.F. 2016. Do molt-migrant songbirds optimize migration routes based on primary productivity? *Behav. Ecol.* **27**: 784–792.
- Contina, A., Bridge, E.S., Seavy, N.S., Duckles, J.M. & Kelly, J.F. 2013. Using geologgers to investigate bimodal isotope patterns in Painted Buntings (*Passerina ciris*). *Auk* **130**: 265–272.
- Contina, A., Harrigan, R.J., Pollinger, J.P., Bridge, E.S., Kelly, J.F. & Smith, T.B. 2016a. Genome-wide identification and characterization of simple sequence repeats in the Painted Bunting (*Passerina ciris*) for population and forensic analyses. *Conserv. Genet. Resour.* **8**: 43–81.
- Contina, A., Bridge, E.S. & Kelly, J.F. 2016b. Exploring novel candidate genes from the Mouse Genome Informatics Database: potential implications for avian migration research. *Integr. Zool.* **11**: 240–249.
- Contina, A., Bay, R., Underwood, V.L., Smith, T.B., Bridge, E.S., Kelly, J.F. & Ruegg, K.C. 2017. Characterization of SNP markers for the Painted Bunting (*Passerina ciris*) and their relevance in population differentiation and genome evolution studies. *Conserv. Genet. Resour.* <https://doi.org/10.1007/s12686-017-0950-7>.
- Contina, A., Bridge, E.S., Ross, J.D., Shipley, J.R. & Kelly, J.F. 2018. Examination of *Clock* and *Adcyap1* gene variation in a neotropical migratory passerine. *PLoS ONE* **13**: e0190859.
- CruzRomo, L. & Oliveras de Ita, A. 2011. Conceptos ecológicos, métodos y técnicas para la conservación y aprovechamiento de aves canoras, de ornato y psitácidos. In Sánchez, O., Zamorano, P., Peters, E. & Moya, H. (eds) *Temas sobre conservación de vertebrados silvestres en México*: 121–148. México, DF: Secretaría de Medio Ambiente y Recursos Naturales, Instituto Nacional de Ecología.
- De Volo, S.B., Reynolds, T., Douglas, M.R. & Antolin, M.F. 2008. An improved extraction method to increase DNA yield from molted feathers. *Condor* **110**: 762–766.
- DePristo, M.A., Banks, E., Poplin, R.E., Garimella, K.V., Maguire, J.R., Hartl, C., Philippakis, A.A., del Angel, G.,

- Rivas, M.A., Hanna, M., McKenna, A., Fennell, T.J., Kernysky, A.M., Sivachenko, A.Y., Cibulskis, K., Gabriel, S.B., Altshuler, D. & Daly, M.J. 2011. A framework for variation discovery and genotyping using next-generation DNA sequencing data. *Nat. Genet.* **43**: 491–498.
- Earl, D.A. & vonHoldt, B.M. 2012. STRUCTURE HARVESTER: a website and program for visualizing STRUCTURE output and implementing the Evanno method. *Conserv. Genet. Resour.* **4**: 359–361.
- Ellison, A.M. 2014. Political borders should not hamper wildlife. *Nature* **508**: 9.
- Evanno, G., Regnaut, S. & Goudet, J. 2005. Detecting the number of clusters of individuals using the software STRUCTURE: a simulation study. *Mol. Ecol.* **14**: 2611–2620.
- Excoffier, L. & Lischer, H.E.L. 2010. Arlequin suite ver 3.5: a new series of programs to perform population genetics analyses under Linux and Windows. *Mol. Ecol. Resour.* **10**: 564–567.
- Francois, O. & Durand, E. 2010. Spatially explicit Bayesian clustering models in population genetics. *Mol. Ecol. Resour.* **10**: 773–784.
- Galimberti, A., Mattia, F.D., Losa, A., Bruni, I., Federici, S., Casiraghi, M., Martellow, S. & Labra, M. 2013. DNA barcoding as a new tool for food traceability. *Food Res. Int.* **50**: 55–63.
- Goudet, J. 2001. FSTAT, a program to estimate and test gene diversity and fixation indices (version 2.9.3). Available at: <http://www2.unil.ch/popgen/softwares/fstat.htm> (accessed September 2017).
- Guo, S. & Thompson, E. 1992. Performing the exact test of Hardy–Weinberg proportion for multiple alleles. *Biometrics* **48**: 361–372.
- Hamilton, R.A. 2001. Records of caged birds in Baja California. In *Monographs in Field Ornithology No. 3*: 254–257. Colorado Springs, CO: American Birding Association.
- Herr, C., Sykes, P. & Klicka, J. 2011. Phylogeography of a vanishing North American songbird: the Painted Bunting (*Passerina ciris*). *Conserv. Genet.* **12**: 1395–1410.
- Howell, S.N.G. & Webb, S. 1995. *A Guide to the Birds of México and Northern Central America*. New York, NY: Oxford University Press.
- Hubisz, M., Falush, D., Stephens, M. & Pritchard, J. 2009. Inferring weak population structure with the assistance of sample group information. *Mol. Ecol. Resour.* **9**: 1322–1332.
- Íñigo-Elias, E.E. 1986. Active trade threatens Mexican avifauna. *TRAFFIC (U.S.A.)* **6**: 6–7.
- Íñigo-Elias, E.E., Rosenberg, K.V. & Wells, J.V. 2002. The Danger of Beauty. *Birdscope*, newsletter Summer 2002 Vol 16, no 3, of the Cornell Lab of Ornithology, Summer 2002. Available at: <http://birds.cornell.edu/publications/birdscope> (accessed September 2017).
- IUCN Species Survival Commission and TRAFFIC 2004. IUCN/TRAFFIC Analyses of the Proposals to Amend the CITES Appendices. Prepared by IUCN Species Survival Commission and TRAFFIC for the Thirteenth Meeting of the Conference of the Parties to CITES. Gland: IUCN – The World Conservation Union.
- Iyengar, A. 2014. Forensic DNA analysis for animal protection and biodiversity conservation: a review. *J. Nat. Conserv.* **22**: 195–205.
- Jakobsson, M. & Rosenberg, N.A. 2007. CLUMPP: a cluster matching and permutation program for dealing with label switching and multimodality in analysis of population structure. *Bioinformatics* **23**: 1801–1806.
- Jonker, R.M., Zhang, Q., Van Hooff, P., Loonen, M.J.J.E., Van der Jeugd, H.P., Crooijmans, R.P., Groenen, M.A., Prins, H.H. & Kraus, R.H. 2012. The development of a genome wide SNP set for the Barnacle Goose *Branta leucopsis*. *PLoS ONE* **7**: e38412.
- Jonzén, N., Lindén, A., Ergon, T., Knudsen, E., Vik, J.O., Rubolini, D., Piacentini, D., Brinch, C., Spina, F., Karlsson, L., Stervander, M., Andersson, A., Waldenström, J., Lehtikoinen, A., Edvardsen, E., Solvang, R. & Stenseth, N.C. 2006. Rapid advance of spring arrival dates in long-distance migratory birds. *Science* **312**: 1959–1961.
- Lopez Medellin, X. & Íñigo-Elias, E.E. 2002. The Passeriformes Cage Trade in Mexico: a Challenge for Bird Conservation. Paper presented at The Third North American Ornithological Conference. New Orleans, 24–28 September 2002.
- Loss, S.R., Will, T. & Marra, P.P. 2013. The impact of free-ranging domestic cats on wildlife of the United States. *Nat. Commun.* **4**: 1396.
- Lowther, P.E., Lanyon, S.M. & Thompson, C.W. 1999. Painted Bunting (*Passerina ciris*). In Poole, A. & Gill, F. (eds) *Birds of North America*, no. 398. Philadelphia, PA: Academy of Natural Sciences and Washington, DC: American Ornithologists' Union.
- Marra, P.P., Hobson, K.A. & Holmes, R. 1998. Linking winter and summer events in a migratory bird by using stable carbon isotopes. *Science* **282**: 1884–1886.
- Marra, P.P., Cohen, E.B., Loss, S.R., Rutter, J.E. & Tonra, C.M. 2015. A call for full annual cycle research in animal ecology. *Biol. Lett.* **11**: 20150552.
- Mlodinow, S.G. & Hamilton, R.A. 2005. Vagrancy of Painted Bunting (*Passerina ciris*) in the United States, Canada, and Bermuda. *N. Am. Birds* **59**: 172–183.
- Moritz, C. 1994. Defining 'evolutionarily significant units' for conservation. *Trends Ecol. Evol.* **9**: 373–375.
- National Audubon Society (NAS) 2004a. Audubon Watchlist Species, Painted Bunting *Passerina ciris*. Available at: <http://audubon2.org/webapp/watchlist/viewSpecies.jsp?id=156> (accessed September 2017).
- National Audubon Society (NAS) 2004b. The Christmas Bird Counts. National Audubon Society. Available at: <http://www.audubon.org/bird/cbc/> (accessed September 2017).
- Nei, M. 1988. *Molecular Evolutionary Genetics*. New York, NY: Columbia University Press.
- Norris, D.R. & Taylor, C.M. 2006. Predicting the consequences of carry-over effects for migratory populations. *Biol. Lett.* **2**: 148–151.
- Ogden, R., Dawnay, N. & McEwing, R. 2009. Wildlife DNA forensics – bridging the gap between conservation genetics and law enforcement. *Endangered Species Res.* **9**: 179–195.
- Paetkau, D., Waits, L.P., Clarkon, P.L., Craighead, L. & Strobeck, C. 1997. An empirical evaluation of genetic distance statistics using microsatellite data from bear (*Ursidae*) populations. *Genetics* **147**: 1943–1957.
- Petit, R.J., El Mousadik, A. & Pons, O. 1998. Identifying populations for conservation on the basis of genetic markers. *Conserv. Biol.* **12**: 844–855.
- Presti, F.T., Guedes, N.M.R., Antas, P.T.Z. & Miyaki, C.Y. 2015. Population genetic structure in Hyacinth Macaws

- (*Anodorhynchus hyacinthinus*) and identification of the probable origin of confiscated individuals. *J. Hered.* **106**: 491–502.
- Pritchard, J.K., Stephens, M. & Donnelly, P.** 2000. Inference of population structure using multilocus genotype data. *Genetics* **155**: 945–959.
- Pyle, P.** 1997. *Identification Guide to North American Birds. Part I. Columbidae to Ploceidae*. Bolinas, CA: Slate Creek Press.
- Pyle, P., Leitner, W.A., Lozano-Angulo, L., Avilez-Teran, F., Swanson, H., Gómez-Limón, E. & Chambers, M.K.** 2009. Temporal, spatial, and annual variation in the occurrence of molt-migrant passerines in the Mexican monsoon region. *Condor* **111**: 583–590.
- Ramos, M.** 1982. *El comercio y la explotación de las aves vivas en México*. Xalapa: INIREB, Cuadernos de Divulgación No. 8.
- Rappole, J.H. & Warner, D.W.** 1980. Ecological aspects of migrant bird behavior in Veracruz, Mexico. In Keast, A. & Morton, E.S. (eds) *Migrant Birds in the Neotropics: Ecology, Behavior, Distribution, and Conservation*: 353–393. Washington, DC: Smithsonian Institution Press.
- Raymond, M. & Rousset, F.** 1995. An exact test for population differentiation. *Evolution* **49**: 1280–1283.
- Raymond, M. & Rousset, F.** 1997. GENEPOP (version 3.1). Updated version of Genepop (version 1.2): population genetics software for exact tests and ecumenism. *J. Heredity* **86**: 248–249.
- Robbins, C.S., Sauer, J.R., Greenberg, R.S. & Droege, S.** 1989. Population declines in North American birds that migrate to the neotropics. *Proc. Natl Acad. Sci. USA* **86**: 7658–7662.
- Rohwer, S., Butler, L.K. & Froehlich, D.R.** 2005. Ecology and demography of east–west differences in molt scheduling of Neotropical migrant passerines. In Greenberg, R. & Marra, P.P. (eds) *Birds of Two Worlds: The Ecology and Evolution of Migratory Birds*: 87–105. Baltimore, MD: Johns Hopkins University Press.
- Rosenberg, N.A.** 2004. DISTRUCT: a program for the graphical display of population structure. *Mol. Ecol. Notes* **4**: 137–138.
- Ruegg, K.C., Anderson, E.C., Paxton, K.L., Apkenas, V., Lao, S., Siegel, R.B., DeSante, D.F., Moore, F. & Smith, T.B.** 2014. Mapping migration in a songbird using high-resolution genetic markers. *Mol. Ecol.* **23**: 5726–5739.
- Rundel, C.W., Wunder, M.B., Alvarado, A.H., Ruegg, K.C., Harrigan, R., Schuh, A., Kelly, J.F., Siegel, R.B., DeSante, D.F., Smith, T.B. & Novembre, J.** 2013. Novel statistical methods for integrating genetic and stable isotopic data to infer individual-level migratory connectivity. *Mol. Ecol.* **22**: 4163–4176.
- Sauer, J.R., Hines, J.E., Gough, G., Thomas, I. & Peterjohn, B.G.** 1997. *The North American Breeding Bird Survey Results and Analysis. Version 96.4*. Laurel: Patuxent Wildlife Research Center. Available at: <http://www.mbr-pwrc.usgs.gov/bbs/bbs96.html> (accessed September 2017).
- Sauer, J.R., Hines, J.E. & Fallon, J.** 2003. *North American Breeding Bird Survey Trend Results (Survey wide). The North American Breeding Bird Survey, Results and Analysis 1966–2002. Version 2003.1*. Laurel: USGS Patuxent Wildlife Research Center.
- Sauer, J.R., Link, W.A., Fallon, J.E., Pardieck, K.L. & Ziolkowski, D.J., Jr** 2013. The North American Breeding Bird Survey 1966–2011: summary analysis and species accounts. *N. Am. Fauna* **79**: 1–32.
- Schuelke, M.** 2000. An economic method for the fluorescent labeling of PCR fragments. *Nat. Biotechnol.* **18**: 233–234.
- Secretaría de Medio Ambiente y Recursos Naturales (SEMARNAT)** 2013. Aprovechamientos. Available at: <http://www.semarnat.gob.mx/temas/gestion-ambiental/vida-silvestre/aprovechamientos> (accessed 10 December 2014).
- Seutin, G., White, B. & Boag, P.** 1991. Preservation of avian blood and tissue samples for DNA analysis. *Can. J. Zool.* **69**: 82–90.
- Sherry, T.W. & Holmes, R.T.** 1993. Are populations of neotropical migrant birds limited in summer or winter? Implications for management. In Finch, D.M. & Stangel, P.W. (eds) *Status and Management of Neotropical Migratory Birds*: 47–57. Fort Collins, CO: U.S. Department of Agriculture.
- Shipley, J.R., Contina, A., Batbayar, N., Bridge, E.S., Peterson, A.T. & Kelly, J.F.** 2013. Niche conservatism and disjunct populations: a case study with Painted Buntings (*Passerina ciris*). *Auk* **130**: 476–486.
- Slatkin, M.** 1995. A measure of population subdivision based on microsatellite allele frequencies. *Genetics* **139**: 457–462.
- Slatkin, M. & Excoffier, L.** 1996. Testing for linkage disequilibrium in genotypic data using the EM algorithm. *Heredity* **76**: 377–383.
- Storer, R.W.** 1951. Variation in the Painted Bunting (*Passerina ciris*), with special reference to wintering populations. *Occ. Papers Mus. Zool. Univ. Michigan* **532**: 1–12.
- Sykes, P.W., Jr & Holzman, S.** 2005. Current range of the eastern population of Painted Bunting (*Passerina ciris*). Part I: breeding. *N. Am. Birds* **59**: 4–17.
- Sykes, P.W., Jr, Holzman, S. & Inigo-Elias, E.E.** 2007. Current range of the eastern population of Painted Bunting (*Passerina ciris*). Part II: winter range. *N. Am. Birds* **61**: 378–406.
- Thompson, C.W.** 1991a. Is the Painted Bunting actually 2 species – problems determining species limits between allopatric populations. *Condor* **93**: 987–1000.
- Thompson, C.W.** 1991b. The sequence of molts and plumages in Painted Buntings and implications for theories of delayed plumage maturation. *Condor* **93**: 209–235.
- U.S. Fish and Wildlife Service (USFWS)** 2004. North Carolina Endangered Species Home Page. Available at: <http://nc-es.fws.gov/es/cntylist/brunswick.html> (accessed September).
- U.S. Fish and Wildlife Service (USFWS)** 2008. *Birds of Conservation Concern*. Arlington, TX: U.S. Department of Interior, Fish and Wildlife Service, Division of Migratory Bird Management. Available at: <http://www.fws.gov/migratorybird/s/> (accessed September).
- Van Oosterhout, C., Hutchinson, W.F., Wills, D.P.M. & Shipley, P.** 2004. MICRO-CHECKER: software for identifying and correcting genotyping errors in microsatellite data. *Mol. Ecol. Notes* **4**: 535–538.
- Waser, P.M. & Strobeck, C.** 1998. Genetic signatures of interpopulation dispersal. *Trends Ecol. Evol.* **13**: 43–44.
- Weir, B.S. & Cockerham, C.C.** 1984. Estimating F-statistics for the analysis of population structure. *Evolution* **38**: 1358–1370.
- Welton, L.J., Siler, C.D., Linkem, C.W., Diesmos, A.C., Diesmos, M.L., Sy, E., Brown, R.M.** 2013a. Dragons in our

midst: phyloforensics of illegally traded Southeast Asian water monitor lizards. *Biol. Cons.* **159**: 7–15.

Wright, S. 1943. Isolation by distance. *Genetics* **28**: 139–156.

Received 18 September 2017;
revision accepted 19 June 2018.

Associate Editor: Jon Martin Collinson.

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section at the end of the article.

Figure S1. STRUCTURE HARVESTER output. The K -values of 2 and 3 are the most likely number of clusters across the Painted Bunting breeding range based on (A) admixture (ADMX) and (B) admixture + LOCPRIOR (ADMX + LOCPRIOR) models, respectively. Model runs were averaged with 10 replicates for each value of K .

Figure S2. (A) Admixture (ADMX) model. (B) Admixture + LOCPRIOR (ADMX + LOCPRIOR) model. Individual Q matrix calculated in STRUCTURE. Each area within two vertical black lines represents a population and the area width is proportional to the number of individuals sampled. The probability (Q) of each individual to be assigned to a single or more clusters is shown on the vertical axis and the clusters are represented by different colours. Model runs averaged with 10 replicates for each value of K .

Table S1. List of sampling latitude and longitude coordinates and population cluster considered for genetic analyses. Sample ID numbers indicated in bold are from Herr *et al.* (2011).

Table S2. List of loci used in the present study and their corresponding code from Contina *et al.* (2016a) and the National Center for Biotechnology Information (NCBI) accession numbers.

Table S3. Expected and observed heterozygosity for each locus for each population and Hardy–Weinberg equilibrium results (significant P -values are highlighted in bold).

Table S4. Schematic table showing linkage disequilibrium results in each population. The + symbol indicates possible occurrence of linked loci.

Table S5. STRUCTURE assignment probabilities for migratory birds collected at the wintering grounds. The probability (Q) of each individual to be assigned to a single clade is shown in the assignment probability columns and ranges from 0 to 1 (with $Q > 0.6$ considered a robust clade assignment value). Assignment probabilities for sample numbers 92, 102, 105 and 106 were rounded to 0.6. The western, central and eastern clades are represented by different colours, and assignments that yielded uncertain results ($Q < 0.6$) are shown in grey.

Table S6. R_{ST} values for pairwise comparisons of populations calculated in ARLEQUIN. Significant values ($P < 0.05$) are highlighted in bold.