

# The prevalence of avian *Plasmodium* is higher in undisturbed tropical forests of Cameroon

Camille Bonneaud<sup>\*,1</sup>, Irem Sepil<sup>\*,2</sup>, Borja Milá<sup>†,3</sup>, Wolfgang Buermann<sup>\*</sup>, John Pollinger<sup>\*</sup>,  
Ravinder N.M. Sehgal<sup>\*,‡</sup>, Gediminas Valkiūnas<sup>§</sup>, Tatjana A. Iezhova<sup>§</sup>,  
Sassan Saatchi<sup>\*,#</sup> and Thomas B. Smith<sup>\*,†</sup>

\* Center for Tropical Research, Institute of the Environment, University of California at Los Angeles, Los Angeles, CA 90095, USA

† Department of Ecology and Evolutionary Biology, University of California at Los Angeles, Los Angeles, CA 90095, USA

‡ Department of Biology, San Francisco State University, San Francisco CA USA

§ Institute of Ecology, Vilnius University, Akademijos 2, LT-08412 Vilnius, Lithuania

# Jet Propulsion Laboratory, California Institute of Technology, Pasadena, CA 94132 USA

(Accepted 16 April 2009)

**Abstract:** Habitat fragmentation and deforestation are thought to disrupt host–parasite interactions and increase the risk of epizootic outbreaks in wild vertebrates. A total of 220 individuals from three species of African rain-forest bird (*Andropadus latirostris*, *Andropadus virens*, *Cyanomitra obscura*), captured in two pristine and two agroforests in Cameroon, were screened for the presence of avian haemosporidian parasites (species of *Plasmodium* and *Haemoproteus*) to test whether habitat differences were associated with differences in the prevalence of infectious diseases in natural populations. Thirteen mitochondrial lineages, including 11 *Plasmodium* and two *Haemoproteus* lineages were identified. Whereas levels of *Haemoproteus* spp. infections were too low to permit analysis, the prevalence of infections with *Plasmodium* spp. reached significantly greater levels in undisturbed mature forests. Importantly however, the significant association between forest type and parasite prevalence was independent of host density effects, suggesting that the association did not reflect changes in host species composition and abundance between forest types. Our results illustrate how characterizing land-cover differences, and hence changes, may be a prerequisite to understanding and predicting patterns of parasite infections in natural populations of rain-forest birds.

**Key Words:** agroforest, *Andropadus latirostris*, *Andropadus virens*, *Cyanomitra obscura*, *Haemoproteus*, host–parasite interaction, *Plasmodium*, primary forest, rain-forest birds

## INTRODUCTION

Human impacts on the environment are increasingly disturbing the ecology of host–parasite relationships, whether through climate change (Kutz *et al.* 2005) or more directly through habitat modifications such as deforestation (Patz *et al.* 2004). Human-driven habitat alterations have even been reported to play a role in the emergence of infectious diseases giving rise to

epidemics (Guerra *et al.* 2006). For example, logging and road building have coincided with an upsurge of leishmaniasis in humans in Latin America (Desjeux 2001), possibly as a result of the adaptation of sand fly vectors to these disturbed habitats (Patz *et al.* 2000). The consequences of these disturbances for wild host–parasite systems are however largely unknown (Gillespie & Chapman 2006, Gillespie *et al.* 2005, Vaz *et al.* 2007) and effects on natural host–parasite interactions may be complex, particularly for vector-borne infections, which involve more than two co-evolving species. When investigating the consequences of forest degradation on the risk of Lyme disease, Allan *et al.* (2003) found that fragmentation increased the density of competent rodent hosts and thereby indirectly the density of infected larval ticks, thus heightening the risks of transmission of the disease to susceptible hosts. Anthropogenic changes of the

<sup>1</sup> Corresponding author. Current address: Department of Organismic and Evolutionary Biology, Museum of Comparative Zoology, Harvard University, USA. Email: cbonneaud@oeb.harvard.edu

<sup>2</sup> Current address: Department of Zoology, The Edward Grey Institute of Field Ornithology, Oxford University, UK

<sup>3</sup> Current address: Museo Nacional de Ciencias Naturales, CSIC, Madrid, Spain

environment may hence threaten wild populations not only through habitat loss, but by perturbing interspecific interactions they may also exacerbate selection acting on natural populations and increase extinction risks. For this reason, it is essential to understand how human-induced habitat differences affect host–parasite dynamics in the wild and define the underlying land-use factors that may act to modify these interactions.

In this study, we explored the habitat specificities of haemosporidian parasites (*Plasmodium* and *Haemoproteus* spp.) in wild bird populations of the rain forests of southern Cameroon. Species of *Plasmodium* and *Haemoproteus* (Haemosporida, Apicomplexa) infect vertebrate hosts and blood-sucking dipteran insects, the vectors of the parasites (Atkinson & Van Riper 1991). The primary vectors of avian *Plasmodium* spp. are mosquitoes belonging among others to the genera *Culex*, *Aedes*, *Culiseta*, *Anopheles* and *Mansonia* (Bequaert 1954, Valkiūnas 2005), whereas *Haemoproteus* spp. are transmitted by blood-sucking biting midges (*Culicoides*) (Kettle 1982, Valkiūnas & Iezhova 2004) and louse-flies (*Pseudolynchia*, *Microlynchia* and *Ornithomyia*) (Bequaert 1954, Kettle 1982, Valkiūnas 2005). Although *Haemoproteus* spp. have traditionally been considered to be relatively host-specific and *Plasmodium* spp. more generalists (Hellgren *et al.* 2007, Ricklefs & Fallon 2002), recent evidence suggests that more or less closely related lineages within both genera may actually exhibit a wide range of host-species with vertebrate-host specificity being frequently broader than host-family level (Beadell *et al.* 2009, Križanauskienė *et al.* 2006). The level of host specificity for haemosporidian parasites may actually depend, in part, on the vector-host association (Gager *et al.* 2008, Hellgren *et al.* 2008).

We tested the hypothesis that habitat differences between pristine and human-altered rain forests were associated with different levels of parasite infections in resident bird populations. Since different bird species may display different levels of infection and because host species composition may vary between mature sites and agroforests, we controlled for changes in host abundance when examining association patterns between forest type and parasite prevalence. We sampled individuals from three rain-forest bird species (*Andropadus latirostris*, Strickland 1844; *Andropadus virens*, Cassin 1857; and *Cyanomitra obscura*, Jardine 1843) from two mature forests and two agroforest sites in southern Cameroon during June and July 2005, just prior to the breeding season. In Cameroon, the amount of mature forest is decreasing, while the proportion of secondary forest is increasing as a result of human activity related to logging and slash-and-burn agriculture (Laurance 1999). Many of the same bird species inhabit these two types of habitat (Louette 1981), yet very little is known about how deforestation can affect parasite prevalence in these populations.

## METHODS

### Sample sites and habitat characterization using remote-sensing data

Fieldwork took place in June and July 2005 at four sites in Cameroon. Locations and dates of fieldwork for the two mature sites are: Zoebefame (2°39'31" N, 13°23'49" E), 24–27 July; Bobo Camp (2°39'17" N, 13°28'16" E), 19–22 July; and for the two agroforest sites: Ndibi (2°43'50" N, 9°52'19" E), 27 June–4 July; Nkwouak (3°52'1" N, 13°18'58" E), 7–12 July. All sites are at similar elevation (600–700 m) and annually undergo two dry seasons and two rainy seasons. The smaller wet season takes place between March and June and is followed by a smaller dry season between June and August, a greater wet season between August and November and finally a greater dry season between November and March. We presented parasite prevalence data from Ndibi from previous years elsewhere (Sehgal *et al.* 2005).

Mature forest sites were located at least 30 km from the nearest road or human settlement and based on ground surveys showed little or no signs of human disturbance. The forest stands were characterized by a layered closed canopy with tall emergent trees and a relatively open understorey. Agroforest sites were adjacent to human settlements and consisted of a mixture of cocoa and coffee trees. These sites had significant disturbance associated with wood harvesting, burning and various other forms of cultivation. The agroforest stands typically had much denser understorey and more broken open canopy than the mature forests. Greater tree cover and vegetation biomass at mature sites were confirmed by remote-sensing observations (for a review see Turner *et al.* 2003). Optical satellite data, expressed as the percentage of tree canopy cover with 500-m spatial resolution, can distinguish open (e.g. shrublands, savannas), fragmented and deforested areas from closed forests (Hansen *et al.* 2002). Microwave remotely sensed data, expressed in decibels (dB), increases with increasing moisture and the overall vegetation biomass (Saatchi *et al.* 2000). Tree canopy cover values (moderate-resolution imaging spectroradiometer surface reflectance of the year 2001, aggregated to 1 km) were: Bobo = 78%, Zoebefame = 81%, Ndibi = 45%, Nkwouak = 68%. Overall vegetation biomass values (Japanese Earth Resource Satellite data in 1996, aggregated to 1 km) were: Bobo = –6.88 dB, Zoebefame = –6.88 dB, Ndibi = –9.70 dB and Nkwouak = –8.85 dB.

### Field methods

At each site, 15–20 mist nets (12 m, 30 × 30-mm mesh) were erected to capture birds. Netting took place generally between daybreak (06h00) and midday (12h00), except

during periods of excessive sun, in which nets were closed earlier. Captured birds were weighed with a 50-g Pesola spring balance, measured, marked with a uniquely numbered aluminium band, and released, following methods described by Smith (1990). Blood samples (1–3 drops) were collected from the brachial vein and stored in lysis buffer (10 mM Tris-HCL pH 8.0, 100 mM EDTA, 2% SDS). We captured 220 birds belonging to the following three species (taxonomy according to Borrow & Demey 2005): yellow-whiskered greenbul (*Andropadus latirostris*, Pycnonotidae,  $n = 71$ ), little greenbul (*Andropadus virens*, Pycnonotidae,  $n = 82$ ) and olive sunbird (*Cyanomitra obscura*, Nectariniidae,  $n = 67$ ). We calculated the relative abundance of each species at each of the four sites by dividing the total number of individuals of a given species captured by the total mist net hours at that site. Mist net hours were calculated as the number of net metres run multiplied by the number of hours nets remained open. Relative abundance is expressed in number of birds  $\times 100/\text{mist-net m} \times \text{h}$ .

All three avian species or related species are common in both mature and agroforest sites and are found throughout equatorial Africa. In a recent study on little greenbul, *A. virens*, Smith *et al.* (2008) found significant differences in morphology, colour plumage and song between individuals captured in mature and secondary forest sites in Cameroon, suggesting that each forest type harbours distinct populations. Populations of *A. virens* can further be distinguished by analysis of amplified fragment length polymorphism, suggesting that, despite detectable gene flow (Smith *et al.* 1997), they remain genetically distinct and individuals rarely move between habitats. This is further corroborated by long-term banding studies suggesting little movement of individuals between habitats (T. B. Smith unpubl. data). Less information is available for *A. latirostris* though long-term banding studies suggest it may be more vagile (Smith unpubl. data). Interestingly, *A. virens* tends to be found at higher densities in secondary than in mature forests, while the reverse is true for *A. latirostris* (Louette & Bijmens 1995). Both bulbul species feed on fruit and insects (Keith *et al.* 1992). The olive sunbird, *C. obscura*, feeds on both nectar and insects and is one of the more common forest sunbirds (Fry *et al.* 1988). The distribution of our three target species across mature and disturbed forests and the relatively restricted movement of individuals between habitats make them good model species for examining the consequences of forest disturbance on natural host–parasite systems.

### Parasite screening using PCR

DNA was extracted from whole blood following a DNeasy kit protocol (Qiagen, Valencia, CA, USA). To

amplify a portion of the cytochrome *b* gene, we used primers L15183 and H15909 and methods described by Szymanski & Lovette (2005). We used 0.25U *Taq* (Qiagen, 201203) in 25  $\mu\text{l}$  volume and ran our amplifications as follows: 50 s at 95 °C, 50 s at 53 °C and 60 s at 72 °C (35 cycles). The PCR products were run on 2% agarose gels and stained with ethidium bromide for UV detection. Negative infections were confirmed by repeated PCR. PCR products were purified using a MinElute Qiagen kit following manufacturer's instructions. We identified lineages by sequencing the fragments (BigDye (R) version 1.1 sequencing kit, Applied Biosystems, Foster City, CA, USA) on an ABI 3730 capillary sequencer (Applied Biosystems). Unresolved sequences showing double peaks in the electropherograms were examined for putative multiple infections by cloning (TOPO-cloning kit, Invitrogen) and sequencing (Pérez-Tris & Bensch 2005). We sequenced 6–10 clones from each sample for which we suspected a multiple infection. We kept only the sequences found several times in independent PCRs, either within the same individual or in several different individuals. We however discarded all unique sequences, which only differed from verified sequences by one nucleotide. Indeed, a single nucleotide divergence may, for example, be attributed to a *Taq* polymerase incorporation error during amplification. All other unique sequences were also discarded.

### Parasite screening using microscopy

We prepared three blood slides for each individual captured. Blood slides were air dried within 5–10 s after their preparation. In a humid environment, we used a battery-operated fan to aid in the drying of the blood films. Slides were fixed in methanol in the field and then stained with Giemsa in the laboratory. The slides were examined for 10–15 min at low magnification ( $\times 400$ ), and then at least 100 fields were studied at high magnification ( $\times 1000$ ). The approximate number of screened red blood cells was  $5 \times 10^5$  in each blood film; the microscopy took approximately 20–25 min per slide. Blood slides were examined with an Olympus BX61 light microscope equipped with an Olympus DP70 digital camera and analysed with the imaging software AnalySIS Five. Parasites were identified according to Valkiūnas (2005). Representative blood slides were deposited in the Institute of Ecology, Vilnius University (Vilnius).

### Phylogenetic and statistical analyses

Sequence divergence was computed using PAUP 4.0b10 (Swofford, D., 2000, PAUP 4.0b7a, Phylogenetic Analysis Using Parsimony (and other methods), Sinauer,

Sunderland, MA). We used Bayesian analysis to construct a phylogeny of parasite cytochrome *b* lineages. We first determined the model of sequence evolution that best fitted the data using MrModeltest (Nylander, J. A. A., 2004, MrModeltest version 2, Evolutionary Biology Centre, Uppsala University, Uppsala, URL <http://www.csit.fsu.edu/~nylander/>). Bayesian analysis of the sequence data was then conducted with MrBayes version 3.1.2 (Huelsenbeck *et al.* 2001) using the model of sequence evolution obtained from MrModeltest (GTR+G), and partitioning sites by codon position. Two Markov chains were run simultaneously for 2 million generations and sampled every 100 generations. We confirmed chain convergence using TRACER 1.3 (Rambaut, D. & Drummond, A. 2003, Tracer v1.3., URL <http://tree.bio.ed.ac.uk/software/tracer/>) and excluded trees generated prior to stationarity. We also generated a phylogeny of haplotypes using the neighbour-joining method and Kimura two-parameter genetic distances as implemented in the program MEGA2 (Kumar *et al.* 2004). Node support for the neighbour-joining tree was determined through 10 000 bootstrap replicates.

The prevalence of infection with *Haemoproteus* spp. was too low to permit analysis. The prevalence of infection with each of the individual *Plasmodium* spp. was similarly prohibitively too low to permit analysis at level of each individual lineage (Table 1). Furthermore, over 70% of all recorded *Plasmodium* spp. infections were closely related malaria parasites belonging to the subgenus *Novyella*, which appears to predominate in African rain-forest passerines (Valkiūnas *et al.* 2009). Consequently, we pooled all infections involving a *Plasmodium* spp. and categorized each bird as either being infected or not. We

used generalized linear mixed models in Genstat (release 9, Rothamsted Experimental Station, Harpenden, UK) with binomial error structure and logit link function to examine: (a) whether bird species differed in infection levels with *Plasmodium* spp. lineages; and (b) whether the relative abundance of host species and forest type was associated with different probabilities of infection with *Plasmodium* spp. In all analyses, site was specified as a random factor to account for a lack of independence of measures within each site.


## RESULTS

We determined haemosporidian parasite prevalence in 220 individuals belonging to three different species of African rain-forest bird. Thirteen verified mitochondrial lineages were found using PCR (eleven *Plasmodium* and two *Haemoproteus* lineages) with numbers of individuals infected ranging from 1 to 20 (Table 1). We conducted BLAST (basic local alignment search tool) analysis of each sequence but found no identical matches of any of the lineages we identified. All sequences are deposited in GenBank (see Figure 1 for the accession numbers). The phylogenetic tree revealed the existence of two monophyletic *Plasmodium* spp. clades of high support (Figure 1). The topologies obtained from Bayesian inference and neighbour joining were almost identical and included only minor topological differences within clades, so only the Bayesian tree is reported on Figure 1. Mean uncorrected sequence divergence between the two *Plasmodium* spp. clades was 0.076 (SD = 0.010), and between *Plasmodium* and *Haemoproteus* lineages was

**Table 1** Total number of birds caught and relative abundance of each bird species expressed in number of birds  $\times$  100/mist-net  $m \times h$ . The three bird species are: *Andropadus latrisrostris* (*A. lat.*), *Andropadus virens* (*A. vir.*) and *Cyanomitra obscura* (*C. obs.*). The number of birds infected by each mitochondrial lineage of *Plasmodium* and *Haemoproteus* spp. detected by PCR is also indicated for each site; PV1, PV2, PV3, PV4, PV6, PV9, PV12, PV13, PV15, PV16, PV17 are the 11 *Plasmodium* spp. lineages and HV1 and HV2 are the two *Haemoproteus* spp. lineages. Bobo and Zobeufame are the two mature forest sites and Ndibi and Nkwouak are the two agroforest sites.

Forest site	Bird species	Total number of birds caught	Relative abundance	PV1	PV2	PV3	PV4	PV6	PV9	PV12	PV13	PV15	PV16	PV17	HV1	HV2
Bobo	<i>A. lat.</i>	18	2.14	5	0	3	0	1	3	0	0	0	0	0	0	0
	<i>A. vir.</i>	6	0.71	0	0	0	0	1	0	0	0	0	0	0	0	0
	<i>C. obs.</i>	21	2.50	0	0	0	0	0	0	0	0	0	0	8	6	3
Zobeufame	<i>A. lat.</i>	32	4.37	2	1	3	0	0	3	0	0	0	0	0	0	0
	<i>A. vir.</i>	8	1.09	1	0	0	0	0	0	0	0	0	0	0	0	0
	<i>C. obs.</i>	19	2.59	0	0	0	0	0	0	0	2	2	1	9	1	0
Ndibi	<i>A. lat.</i>	13	1.38	1	2	0	1	0	2	0	0	0	0	0	0	0
	<i>A. vir.</i>	38	4.02	2	0	0	0	0	1	0	0	0	0	0	0	0
	<i>C. obs.</i>	7	0.74	0	0	0	0	0	0	0	0	2	0	1	0	1
Nkwouak	<i>A. lat.</i>	8	1.03	0	0	0	0	0	1	1	0	0	0	0	0	0
	<i>A. vir.</i>	30	3.86	1	0	0	0	0	0	0	0	0	0	0	0	0
	<i>C. obs.</i>	20	2.57	0	0	0	0	0	0	0	0	2	2	2	0	1
Total		220		12	3	6	1	2	10	1	2	6	3	20	7	5






**Figure 1.** Phylogenetic relationships among haplotypes from the 13 malaria parasites (PCR detection) found in three species of rain-forest bird in Cameroon (*A. lat.*: *Andropadus latirostris*, *A. vir.*: *Andropadus virens* and *C. obs.*: *Cyanomitra obscura*), based on Bayesian analysis of *Cyt b* sequences. The tree corresponds to the consensus topology of 30 002 trees obtained after 2 million generations based on the GTR+G model of sequence evolution. A sequence from *Leucocytosoon sabrazesi* (Mathis & Léger 1910) (AB299369) was used as outgroup. We also added a sequence from *Haemoproteus belopolskyi* (Valkiūnas 1989) (DQ630006). The two highly supported monophyletic *Plasmodium* spp. clades are indicated on the figure. GenBank accession numbers of all sequences are indicated. Numbers along branches correspond to node support from Bayesian analysis (left) and bootstrap analysis of the neighbour-joining reconstruction (right). The presence of each parasite lineage in the three species of rain-forest birds is indicated by black solid fill. PV9 corresponds to the morphospecies *Plasmodium multivacuolaris*, PV15 to *Plasmodium megaglobularis* and PV17 to *Plasmodium lucens* (Valkiūnas *et al.* 2008b, 2009).

0.125 (SD = 0.015). The two monophyletic *Plasmodium* spp. clades displayed interesting host-genus specificity, since parasites of clade 1 infected only the *Andropadus* species and parasites of clade 2 infected mostly *C. obscura* (Figure 1). Indeed, in the latter case, only PV12 was found to infect *A. latirostris*. *Haemoproteus* lineages were found only in *C. obscura* and the prevalence of the infection was too low to test for any association with habitat characteristics. Interestingly, the microscopy approach allowed us to link three *Plasmodium* spp. lineages identified using PCR with morphospecies belonging to the subgenus *Novyella*: PV9 corresponded to *Plasmodium multivacuolaris*, PV15 to *Plasmodium megaglobularis* and PV17 to *Plasmodium lucens* (Valkiūnas *et al.* 2008b, 2009).

Bird species displayed significantly different levels of *Plasmodium* spp. infections, both when considering PCR data (generalized linear mixed model,  $\chi^2 = 22.9$ ,  $P < 0.001$ ; mean percentage infected individuals predicted by the model,  $N_{A. latirostris} = 40\%$ ,  $N_{A. virens} = 8\%$ ,  $N_{C. obscura} = 47\%$ ) and microscopy data (generalized linear mixed model,  $\chi^2 = 23.8$ ,  $P < 0.001$ ; predicted means:  $N_{A. latirostris} = 51\%$ ,  $N_{A. virens} = 15\%$ ,  $N_{C. obscura} = 52\%$ ). Forest type, however, was significantly associated with differences in the prevalence of *Plasmodium* parasites, both when using PCR data (generalized linear mixed model, host relative abundance:  $\chi^2 = 8.72$ ,  $P = 0.003$ , forest type:  $\chi^2 = 16.0$ ,  $P = < 0.001$ , Figure 2) and microscopy data (generalized linear mixed model, host relative abundance:  $\chi^2 = 2.34$ ,  $P = 0.13$ , forest type:  $\chi^2 = 12.47$ ,  $P < 0.001$ , Figure 2). In both cases, birds were more likely to be infected in mature forest sites than in agroforest sites.



**Figure 2.** Probability that the birds were infected with *Plasmodium* parasites for both habitat types (mature forests and agroforests) modelled using a generalized linear mixed model with site as a random factor. Figures show back-transformed mean probabilities of infection ( $\pm$  SE) for *Plasmodium* spp. detected by PCR and microscopy.

## DISCUSSION

Undisturbed tropical forests were found to display significantly higher prevalence of infection with avian *Plasmodium* spp. (all lineages pooled), both when using PCR and microscopy methods of detection. The discrepancies in the prevalence of haemosporidian blood parasites that we detected when using microscopy and PCR methods are not surprising and have been reported in previous studies (Krone *et al.* 2008, Pérez-Tris *et al.* 2007, Valkiūnas *et al.* 2006a). This highlights the benefit of using a combination of both methods to evaluate the prevalence of haemosporidian blood parasites (Hellgren *et al.* 2007, Križanauskienė *et al.* 2006, Valkiūnas *et al.* 2008a). In any case, our results suggest that forest density and structure may influence interactions between species and play a role in the transmission and/or maintenance of infections.

In a recent study, Wood *et al.* (2007) detected a correlation between small-scale spatial variation determined by GIS analysis, and differences in malaria prevalence in a temperate woodland population of blue tits (*Cyanistes caeruleus*). Parasite prevalence significantly fluctuated by up to six-fold over 1 km, suggesting that individuals of a same population may, in fact, be exposed to considerable variation in the risk of infection. The likely role of landscape in heterogeneous disease dynamics highlights the need to incorporate habitat structure in ecological and epidemiological studies of natural host–parasite systems (Wood *et al.* 2007). An interesting difficulty is to estimate the most relevant scale at which habitat characteristics should be incorporated into assessments of risk or incidence of infection (Ostfeld *et al.* 2005), especially as spatial structure should differentially impact species with varying habitat range and dispersal abilities.

Forest type was associated with differences in the prevalence of *Plasmodium* spp. after we removed the effects of changes in the relative abundance of avian hosts, suggesting that habitat effects on infection patterns do not solely result from changes in host species composition between mature and agroforests. Changes in tropical habitats have been previously reported to correlate with changes in species interactions that were not merely attributable to differences in species richness (Tylianalis *et al.* 2007). Furthermore, correlations between habitat characteristics and parasite prevalence independent of host-density effects have been reported elsewhere. For example, Gillespie & Chapman (2006) detected a significant influence of habitat degradation on the prevalence of gastrointestinal parasites recovered from primate faeces that was not driven by changes in host density. In any case, the pattern detected in this study is actually not surprising in the case of arthropod-borne diseases, as habitat effects may also be mediated via

effects on vector populations. Our results do not however preclude any indirect effects of host abundance, especially as host density is known to positively correlate with parasite prevalence (Morand & Poulin 1998, Packer *et al.* 1999). Infection levels may, in fact, also depend on changes in the abundance of other susceptible host species. Larger screenings of parasite infections in rain-forest birds of Cameroon that were conducted in parallel with this study, established that six of the *Plasmodium* spp. lineages described here were recorded only in certain bird species or bird families, suggesting that they may actually display stringent host-species or host-family specificity (PV1, PV2, PV6, PV9, PV16, PV17) (unpubl. data). These screenings also demonstrated that the other five lineages were found to infect several bird families and hence characterized as generalists (unpubl. data). Because we pooled all *Plasmodium* spp. lineages due to small sample sizes, it is possible that habitat effects on other avian hosts may, to a certain extent, be driving changes in parasite prevalence between forest types.

Effects of habitat on parasite prevalence may be explained by five non-mutually exclusive hypotheses; the predictions from which can be tested in future studies. First, the lower levels of infection with *Plasmodium* spp. in forests of lower vegetation biomass may be explained in part by the fitness cost associated with *Plasmodium* spp. infections (Merino *et al.* 2000, Valkiūnas *et al.* 2006b). Higher vegetation density, especially in times of drought, may indeed protect weakened birds from increased predation risk and thereby increase the number of malaria-infected individuals at mature forest sites. Second, since vegetation is also thought to be crucial in providing cover for dipteran breeding sites (Mercer *et al.* 2005), it is possible that mature forests also offer better vector breeding grounds than agroforests.

Third, increased competition among insect vectors in agroforests may cause a decline in the abundance of vectors susceptible to avian *Plasmodium* spp., resulting in a decreased prevalence of infection. Recent studies have demonstrated that the mosquito *Anopheles darlingi*, which is the primary malaria vector for humans in the Amazon, was encountered in altered habitats but was absent from intact sites (Tadei *et al.* 1998), and that biting rates increased by almost 300% in disturbed habitats (Vittor *et al.* 2006). Furthermore, larval studies suggest that the increase in *A. darlingi* abundance is due to an increase in suitable breeding sites in secondary forests (Vittor *et al.* 2006). The transmission cycles of native pathogens have been shown to be altered by such new vectors (Juliano & Lounibos 2005). If lower prevalence in agroforests results, at least in part, from an intensification of vector competition, then we would expect breeding sites in agroforests to favour vectors preferentially feeding on humans at the detriment of those preferentially feeding on wild birds.

Fourth, the decreased prevalence of infection in agroforests could also be explained by changes in the feeding habits of insect vectors, especially in vectors that are relatively non-selective in their food preferences (Walsh *et al.* 1993). In fact, both agroforest sites (Ndibi and Nkwouak) were close to human settlements (~0.5 km) and human-biting rates of mosquito increased noticeably at those sites compared with mature forest sites (pers. obs.). As a result, mosquitoes may preferentially feed on novel hosts in agroforest sites, such as humans or other animal species (e.g. domestic animals), thereby decreasing the risk of disease transmission to wild birds. This could be tested through the molecular identification of vertebrate hosts in dipteran blood meals. Last, there may be an increase in the density of alternative hosts in agroforest, which are less susceptible to infection (or incompetent). This may consequently result in a decrease in the prevalence and in the risk of infection through a 'dilution effect' (Ostfeld & Keesing 2000, Schmidt & Ostfeld 2001, Yahner 1988).

Here we show that forest type may have notable effects on host–parasite interactions. Our system is particularly complex because it involves many players: several species of avian host, several species and even genera of parasites and dipteran vectors. To fully understand the impact of habitat differences and modifications on these coevolutionary interactions, future efforts will focus on intense sampling of vertebrates (avian, mammalian and reptilian) and dipterans diversity and abundance in order to identify the potential changes in vector abundance and in vector meal preference. Further research will also be required to more fully examine the influence of spatial variation on parasite–host interactions. A weakness of the present study is that only four sites were compared and the two mature forest sites were adjacent. Nevertheless, our preliminary results emphasize the need to incorporate environmental abiotic and biotic characteristics when examining host–parasite dynamics or, more generally, when examining interactions among species. Furthermore, they stress the need to take into account both species abundance and complex species interactions when evaluating the ecological impacts of habitat degradation and planning effective conservation actions. This is particularly imperative given the increasing trend in emerging infectious diseases (Jones *et al.* 2008).

## ACKNOWLEDGEMENTS

We thank the Government of the Republic of Cameroon for permission to conduct the field research. We thank D. Anye, T. Chaser and T. Dietsch for assistance in the field. We also thank R. Ricklefs, two anonymous referees and the Editor for their insightful comments on a previous

version of the manuscript. The research was funded by the joint NSF-NIH Ecology of Infectious Diseases Program award EF-0430146 and by the Lithuanian State Science and Studies Foundation.

## LITERATURE CITED

- ALLAN, B. F., KEESING, F. & OSTFELD, R. S. 2003. Effect of forest fragmentation on Lyme disease risk. *Conservation Biology* 17:267–272.
- ATKINSON, C. T. & VAN RIPER, C. 1991. Pathogenicity and epizootiology of avian haematozoa: *Plasmodium*, *Leucocytozoon*, and *Haemoproteus*. Pp. 19–48 in Loye, E. & Zuk, M. (eds.). *Bird–parasite interactions: ecology, evolution, and behavior*. Oxford University Press, New York.
- BEADELL, J. S., COVAS, R., GEBHARD, C., ISHTIAQ, F., MELO, M., SCHMIDT, B. K., PERKINS, S. L., GRAVES, G. R. & FLEISCHER, R. C. 2009. Host associations and evolutionary relationships of avian blood parasites from West Africa. *International Journal for Parasitology* 39:257–266.
- BEQUAERT, J. C. 1954. The Hippoboscidae or louse-flies (Diptera) of mammals and birds. Part II. Taxonomy, evolution and revision of American genera and species. *Entomologica Americana* 34:1–232.
- BORROW, N. & DEMEY, R. 2005. *Birds of Western Africa*. Princeton Field Guides, New Jersey. 512 pp.
- DESJEUX, P. 2001. The increase in risk factors for leishmaniasis worldwide. *Transactions of the Royal Society of Tropical Medicine and Hygiene* 95:239–243.
- FRY, C. H., KEITH, S. & URBAN, E. K. 1988. *The birds of Africa*. Academic Press, New York. 611 pp.
- GAGER, A. B., LOAIZA, J. D. R., DEARBORN, D. C. & BERMINGHAM, E. 2008. Do mosquitoes filter the access of *Plasmodium* cytochrome b lineages to an avian host? *Molecular Ecology* 17:2552–2561.
- GILLESPIE, T. R. & CHAPMAN, C. A. 2006. Prediction of parasite infection dynamics in primate metapopulations based on attributes of forest fragmentation. *Conservation Biology* 20:441–448.
- GILLESPIE, T. R., CHAPMAN, C. A. & GREINER, E. C. 2005. Effects of logging on gastrointestinal parasite infections and infection risk in African primates. *Journal of Applied Ecology* 42:699–707.
- GUERRA, C. A., SNOW, R. W. & HAY, S. I. 2006. A global assessment of closed forests, deforestation and malaria risk. *Annals of Tropical Medicine and Parasitology* 100:189–204.
- HANSEN, M. C., DEFRIES, R. S., TOWNSHEND, J. R. G., SOHLBERG, R. A., DIMICELI, C. & CARROLL, M. 2002. Towards an operational MODIS continuous field of percent tree cover algorithm: examples using AVHRR and MODIS data. *Remote Sensing of the Environment* 83:303–319.
- HELLGREN, O., WALDENSTROM, J., PEREZ-TRIS, J., SZOLLOSI, E., HASSELQUIST, D., KRÍŽANAUSKIENĖ, A., OTTOSSON, U. & BENSCH, S. 2007. Detecting shifts of transmission areas in avian blood parasites – a phylogenetic approach. *Molecular Ecology* 16:1281–1290.
- HELLGREN, O., BENSCH, S. & MALMQVIST, B. 2008. Bird hosts, blood parasites and their vectors – associations uncovered by molecular analyses of blackfly blood meals. *Molecular Ecology* 17:1605–1613.

- HUELSENBECK, J. P., RONQUIST, F., NIELSEN, R. & BOLIBACK, J. P. 2001. Bayesian inference of phylogeny and its impact on evolutionary biology. *Science* 294:2310–2314.
- JONES, K. E., PATEL, N. G., LEVY, M. A., STOREYGARD, A., BALK, D., GITTLEMAN, J. L. & DASZAK, P. 2008. Global trends in emerging infectious diseases. *Nature* 451:990–993.
- JULIANO, S. A. & LOUNIBOS, L. P. 2005. Ecology of invasive mosquitoes: effect on resident species and on human health. *Ecological Letters* 8:558–574.
- KEITH, S., URBAN, E. K. & FRY, C. H. 1992. *The birds of Africa*. Academic Press, New York. 609+ pp.
- KETTLE, D. S. 1982. *Medical and veterinary entomology*. Wiley and Sons, New York. 658 pp.
- KRIŽANAUSKIENĖ, A., HELLGREN, O., KOSAREV, V., SOKOLOV, L., BENSCH, S. & VALKIŪNAS, G. 2006. Variation in host specificity between species of avian hemosporidian parasites: evidence from parasite morphology and cytochrome B gene sequences. *Journal of Parasitology* 92:1319–1324.
- KRONE, O., WALDENSTROM, J., VALKIUNAS, G., LESSOW, O., MULLER, K., IEZHOVA, T. A., FICKEL, J. & BENSCH, S. 2008. Haemosporidian blood parasites in European birds of prey and owls. *Journal of Parasitology* 94:709–715.
- KUMAR, S., TAMURA, K. & NEI, M. 2004. MEGA3: integrated software for molecular evolutionary genetics analysis and sequence alignment. *Briefings in Bioinformatics* 5:150–163.
- KUTZ, S. J., HOBERG, E. P., POLLEY, L. & JENKINS, E. J. 2005. Global warming is changing the dynamics of Arctic host–parasite systems. *Proceedings of the Royal Society B-Biological Sciences* 272:2571–2576.
- LAURANCE, W. F. 1999. Reflections on the tropical deforestation crisis. *Biological Conservation* 91:109–117.
- LOUETTE, M. 1981. The birds of Cameroon. An annotated checklist. *Verhandelingen van de Koninklijke Academie voor Wetenschappen, Letteren Schone Kunsten van België. Klasse de Wetenschappen* 43:1–295.
- LOUETTE, M. & BIJNENS, L. 1995. The utility of birds as bioindicators – case-studies in Equatorial Africa. *Belgian Journal of Zoology* 125:157–165.
- MERCER, D. R., SHEELEY, S. L. & BROWN, E. J. 2005. Mosquito (Diptera: Culicidae) development within microhabitats of an Iowa wetland. *Journal of Medical Entomology* 42:685–693.
- MERINO, S., MORENO, J., SANZ, J. J. & ARRIERO, E. 2000. Are avian blood parasites pathogenic in the wild? A medication experiment in blue tits (*Parus caeruleus*). *Proceedings of the Royal Society of London Series B, Biological Sciences* 267:2507–2510.
- MORAND, S. & POULIN, R. 1998. Density, body mass and parasite species richness of terrestrial mammals. *Evolutionary Ecology* 12:717–727.
- OSTFELD, R. S. & KEESING, F. 2000. Biodiversity and disease risk: the case of Lyme disease. *Conservation Biology* 14:722–728.
- OSTFELD, R. S., GLASS, G. & KEESING, F. 2005. Spatial epidemiology: an emerging (or re-emerging) discipline. *Trends in Ecology and Evolution* 20:328–336.
- PACKER, C., ALTIZER, S., APPEL, M., BROWN, E., MARTENSON, J., O'BRIEN, S. J., ROELKE-PARKER, M., HOFMANN-LEHMANN, R. & LUTZ, H. 1999. Viruses of the Serengeti: patterns of infection and mortality in African lions. *Journal of Animal Ecology* 68:1161–1178.
- PATZ, J. A., GRACZYK, T. K., GELLER, N. & VITTOR, A. Y. 2000. Effects of environmental change on emerging parasitic diseases. *International Journal of Parasitology* 30:1395–1405.
- PATZ, J. A., DASZAK, P., TABOR, G. M., AGUIRRE, A. A., PEARL, M., EPSTEIN, J., WOLFE, N. D., KILPATRICK, A. M., FOUFOPOULOS, J., MOLYNEUX, D. & BRADLEY, D. J. 2004. Unhealthy landscapes: policy recommendations on land use change and infectious disease emergence. *Environmental Health Perspectives* 112:1092–1098.
- PÉREZ-TRIS, J. & BENSCH, S. 2005. Diagnosing genetically diverse avian malarial infections using mixed-sequence analysis and TA-cloning. *Parasitology* 131:15–23.
- PÉREZ-TRIS, J., HELLGREN, O., KRIŽANAUSKIENĖ, A., WALDENSTROM, J., SECONDI, J., BONNEAUD, C., FJELDSÅ, J., HASSELQUIST, D. & BENSCH, S. 2007. Within-host speciation of malaria parasites. *PLoS ONE* 2:e235.
- RICKLEFS, R. E. & FALLON, S. M. 2002. Diversification and host switching in avian malaria parasites. *Proceedings of the Royal Society of London Series B, Biological Sciences* 269:885–892.
- SAATCHI, S., NELSON, B., PODEST, E. & HOLT, J. 2000. Mapping land cover types in the Amazon Basin using 1 km JERS-1 mosaic. *International Journal of Remote Sensing* 21:1201–1234.
- SCHMIDT, K. A. & OSTFELD, R. S. 2001. Biodiversity and the dilution effect in disease ecology. *Ecology* 82:609–619.
- SEHGAL, R. N. M., JONES, H. I. & SMITH, T. B. 2005. Blood parasites of some West African rainforest birds. *Journal of Veterinary Medical Science* 67:295–301.
- SMITH, T. B. 1990. Resource use by bill morphs of an African finch: evidence for intraspecific competition. *Ecology* 71:1246–1257.
- SMITH, T. B., WAYNE, R. K., GIRMAN, D. J. & BRUFORD, M. W. 1997. A role for ecotones in generating rainforest biodiversity. *Science* 276:1855–1857.
- SMITH, T. B., MILÁ, B., GREETHER, G. F., SLABBEKOORN, H., SEPIL, I., BUERMANN, W., SAATCHI, S. & POLLINGER, J. P. 2008. Evolutionary consequences of human disturbance in a rainforest bird species from central Africa. *Molecular Ecology* 17:58–71.
- SZYMANSKI, M. M. & LOVETTE, I. J. 2005. High lineage diversity and host sharing of malarial parasites in a local avian assemblage. *Journal of Parasitology* 91:768–774.
- TADEI, W. P., THATCHER, B. D., SANTOS, J. M. H., SCARPASSA, V. M., RODRIGUES, I. B. & RAFAEL, M. S. 1998. Ecologic observations on anopheline vectors of malaria in the Brazilian Amazon. *American Journal of Tropical Medicine and Hygiene* 59:325–335.
- TURNER, W., SPECTOR, S., GARDINER, N., FLADELAND, M., STERLING, E. & STEININGER, M. 2003. Remote sensing for biodiversity science and conservation. *Trends in Ecology and Evolution* 18:306–314.
- TYLIANALIS, J. M., TSCHARNTKE, T. & LEWIS, O. T. 2007. Habitat modification alters the structure of tropical host–parasitoid food webs. *Nature* 445:202–205.
- VALKIŪNAS, G. 2005. *Avian malaria parasites and other haemosporidia*. CRC Press, Boca Raton. 932 pp.
- VALKIŪNAS, G. & IEZHOVA, T. A. 2004. The transmission of *Haemoproteus belopoluskyi* (Haemosporida: Haemoproteidae) of blackcap by *Culicoides impunctatus* (Diptera: Ceratopogonidae). *Journal of Parasitology* 90:196–198.



- VALKIŪNAS, G., BENSCH, S., IEZHOVA, T. A., KRIŽANAUSKIENĖ, A., HELLGREN, O. & BOLSHAKOV, C. V. 2006a. Nested cytochrome B polymerase chain reaction diagnostics underestimate mixed infections of avian blood haemosporidian parasites: microscopy is still essential. *Journal of Parasitology* 92:418–422.
- VALKIŪNAS, G., ŽIČKUS, T., SHAPOVAL, A. P. & IEZHOVA, T. A. 2006b. Effect of *Haemoproteus belopolskyi* (Haemosporida: Haemoproteidae) on body mass of the blackcap *Sylvia atricapilla*. *Journal of Parasitology* 92:1123–1125.
- VALKIŪNAS, G., IEZHOVA, T. A., KRIŽANAUSKIENĖ, A., PALINAUSKAS, V., SEHGAL, R. N. M. & BENSCH, S. 2008a. A comparative analysis of microscopy and PCR-based detection methods for blood parasites. *Journal of Parasitology* 94:1395–1401.
- VALKIŪNAS, G., IEZHOVA, T. A., LOISEAU, C., CHASAR, A., SMITH, T. B. & SEHGAL, R. N. M. 2008b. New species of haemosporidian parasites (Haemosporida) from African rainforest birds, with remarks on their classification. *Parasitology Research* 103:1213–1228.
- VALKIŪNAS, G., IEZHOVA, T. A., LOISEAU, C., CHASAR, A., SMITH, T. B. & SEHGAL, R. N. M. 2009. New malaria parasites of the subgenus *Novyella* in African rainforest birds, with remarks on their high prevalence, classification and diagnostics. *Parasitology Research* 104:1061–1077.
- VAZ, V. C., D'ANDREA, P. S. & JANSEN, A. M. 2007. Effects of habitat fragmentation on wild mammal infection by *Trypanosoma cruzi*. *Parasitology* 134:1785–1793.
- VITTOR, A. Y., GILMAN, R. H., TIELSCH, J., GLASS, G., SHIELDS, T., LOZANO, W. S., PINEDO-CANCINO, V. & PATZ, J. A. 2006. The effect of deforestation on the human-biting rate of *Anopheles darlingi*, the primary vector of *falciparum* malaria in the Peruvian Amazon. *American Journal of Tropical Medicine and Hygiene* 74:3–11.
- WALSH, J. F., MOLYNEAUX, D. H. & BIRLEY, M. H. 1993. Deforestation: effects on vector-borne disease. *Parasitology* 106:S55–S75.
- WOOD, M. J., COSGROVE, C. L., WILKIN, T. A., KNOWLES, S. C. L., DAY, K. P. & SHELDON, B. C. 2007. Within-population variation in prevalence and lineage distribution of avian malaria in blue tits, *Cyanistes caeruleus*. *Molecular Ecology* 16:3263–3273.
- YAHNER, R. H. 1988. Changes in wildlife communities near edges. *Conservation Biology* 2:333–339.